

Climate Change and the Kidney

Richard J. Johnson^a Laura G. Sánchez-Lozada^b Lee S. Newman^c
Miguel A. Lanaspá^a Henry F. Diaz^d Jay Lemery^e
Bernardo Rodriguez-Iturbe^f Dean R. Tolan^g Jaime Butler-Dawson^c
Yuka Sato^a Gabriela Garcia^a Ana Andres Hernando^a
Carlos A. Roncal-Jimenez^a

^aDivision of Renal Diseases and Hypertension, University of Colorado, Aurora, CO, USA; ^bDepartment of Nephrology, INC Ignacio Chavez, Mexico City, Mexico; ^cDepartment of Environmental and Occupational Health, Center for Health, Work, and Environment, Colorado School of Public Health, University of Colorado, Aurora, CO, USA; ^dDepartment of Geography and Environment, University of Hawai'i at Mānoa, Honolulu, HI, USA; ^eDivision of Wilderness Health, Department of Emergency Medicine, University of Colorado, Aurora, CO, USA; ^fUniversidad del Zulia, Maracaibo, Zulia, Venezuela; ^gDepartment of Biochemistry, Boston University, Boston, MA, USA

Keywords

Global warming · Heatstroke · Heat waves · Mesoamerican nephropathy · Kidney stones

Abstract

The worldwide increase in temperature has resulted in a marked increase in heat waves (heat extremes) that carries a markedly increased risk for morbidity and mortality. The kidney has a unique role not only in protecting the host from heat and dehydration but also is an important site of heat-associated disease. Here we review the potential impact of global warming and heat extremes on kidney diseases. High temperatures can result in increased core temperatures, dehydration, and blood hyperosmolality. Heatstroke (both clinical and subclinical whole-body hyperthermia) may have a major role in causing both acute kidney disease, leading to increased risk of acute kidney injury from rhabdomyolysis, or heat-induced inflammatory injury to the kidney. Recurrent heat and dehydration can result in chronic kidney disease (CKD) in animals and theoretically plays a role in epidemics of CKD developing in hot regions of the world where workers

are exposed to extreme heat. Heat stress and dehydration also has a role in kidney stone formation, and poor hydration habits may increase the risk for recurrent urinary tract infections. The resultant social and economic consequences include disability and loss of productivity and employment. Given the rise in world temperatures, there is a major need to better understand how heat stress can induce kidney disease, how best to provide adequate hydration, and ways to reduce the negative effects of chronic heat exposure.

© 2019 The Author(s)
Published by S. Karger AG, Basel

Introduction

Increasing worldwide temperatures are now well documented, and the mean temperature increase in the last 50 years approximates 0.8 °Centigrade. While the absolute rise in temperature may not seem large, it is already having major effects on human health [1]. One of the more striking consequences is a marked increase in extreme heat events, termed heat waves [2–4]. Heat waves are the

KARGER

E-Mail karger@karger.com
www.karger.com/anm


© 2019 The Author(s)
Published by S. Karger AG, Basel

Karger
Open access

This article is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND) (<http://www.karger.com/Services/OpenAccessLicense>). Usage and distribution for commercial purposes as well as any distribution of modified material requires written permission.

Richard J. Johnson, MD
Division of Renal Diseases and Hypertension, 12700 East 19th Ave
Room 7012, University of Colorado Anschutz Medical Campus
Aurora, CO 80045 (USA)
E-Mail Richard.Johnson@ucdenver.edu

Fig. 1. Change in heat wave exposure (in individuals >65 years old) relative to the 1986–2008 average. Here heat waves are defined as a period of 3 days or more in which the minimal temperature exceeds the 99th percentile of the average temperature between 1986 and 2008, and the population was limited to individuals 65 years or older. Reused with permission from the Lancet [7].



most common cause of mortality of all weather-related events in the United States (including tornados, hurricanes, and lightning strikes) [5]. Heat waves are also among the top 10 worldwide causes of death by natural disasters between 1980 and 2017 (Table 1). Different definitions have been used to classify heat waves, but one of the more common definitions is a temperature that is 5°C greater than the mean high temperature for a given day, and one that persists for at least 5 days [6]. Numerous studies have reported a dramatic increase in heat waves worldwide [7] (Fig. 1). For example, one estimate suggests that in 2015 alone there were 175 million more people exposed to heat waves as a consequence of climate change [7]. Heat waves are dangerous not only because of the risk of overheating the human body but also due to increased mortality of individuals with cardiovascular and respiratory disorders [6]. One of the worse heat waves was the one that struck Europe in August 2003, resulting in 73,000 deaths [8]. However, heat waves have caused significant mortality throughout the world, including Chicago in 1995 [9], Andhra Pradesh in 2014 and 2015 [10, 11], and Karachi, Pakistan in 2015 [12]. Heat waves and extreme heat also affect labor performance and efficiency [13] and may affect crop performance [14]. Some heat waves have been associated with such extreme temperatures that even wildlife are endangered, and there are predictions that in the future some parts of the world could become so hot that they will become uninhabitable [15–17]. Thus, regulating body temperature is a key to survival. Maintaining a well-hydrated state is critical to this process, but it is of additional concern that the availability of safe water supplies is dwindling worldwide. Indeed, there is now evidence that as much as 10% of the world population faces

Table 1. The most fatal natural disasters between 1980 and 2017

Event (location, date)	Deaths
Tsunami/Earthquake (Thailand, 2004)	220,000
Earthquake (Haiti, 2010)	159,000
Cyclone, storm surge (Myanmar, 2008)	140,000
Cyclone, storm Surge (Bangladesh, 1991)	139,000
Earthquake (Pakistan, 2005)	88,000
Earthquake (China, 2008)	84,000
Heat wave, Drought (Europe, 2003)	70,000
Heat wave (Russia, 2010)	56,000
Earthquake (Iran, 1990)	40,000
Earthquake (Iran, 2003)	26,000

Adapted from data published by Münchener Rückversicherungs-Gesellschaft. Heat waves accounted for the 7th and 8th most fatal natural disasters during this period. From: https://www.munichre.com/site/corporate/get/params_E1716525033_Dattachment/1707976/munichre-natural-catastrophes-in-2018.pdf.

a serious shortage of water availability [18, 19]. In addition, studies suggest that many individuals, including children and adolescents, who do have access to potable water are considered to be underhydrated [20].

In this review, we discuss the effect of climate change on diseases of the kidney. The kidneys have a supreme function in maintaining blood volume to support blood pressure as well as extracellular and intracellular osmolality (“the internal milieu”) that allows for normal metabolism. One of their more important functions is urinary concentration, in which it minimizes fluid loss while assuring the excretion of nitrogenous wastes. Unfortunately, the high metabolic work, as well as the concentrated

excretion of wastes, makes the kidney very susceptible to injury from climate change. Indeed, studies have reported that increasing temperatures translate into increased admissions through the emergency room of a wide range of renal disorders, including acute kidney injury, chronic kidney disease (CKD), kidney stones, and urinary tract infections (UTIs) [21, 22]. Occupational exposure to heat stress has also been linked with higher incidence rates of kidney disease [23] and to a loss in productivity of workers when their kidney function becomes compromised [13]. Here we discuss some of these associations as well as potential links with the epidemics of CKD of unknown origin in hot regions throughout the world.

Heatstroke and Acute Kidney Injury

One of the major health consequences of extreme heat is heatstroke, which results when one cannot adequately control body temperature, resulting in hyperthermia (typically defined as a temperature $>40.6^{\circ}\text{C}$, $>105^{\circ}\text{F}$) that can lead to delirium, coma, seizures, and multiorgan failure [5]. Heatstroke can occur during heat waves (termed epidemic or classic heatstroke) and also in association with exercise or labor in the heat (termed exertional heatstroke) [24].

Exertional heatstroke is especially common among military personnel, marathon runners [25–28], as well as workers in mines or agricultural fields (especially sugarcane) [29]. It is especially common among new workers who are not acclimatized and those that are overweight [29]. Epidemic heatstroke most commonly occurs in association with heat waves and affects those vulnerable to illness, such as the elderly, those with obesity or diabetes, those who are malnourished, individuals who have no air conditioning, and those with underlying cardiovascular or respiratory diseases.

Both classical and exertional heatstroke can be severe, in which case they are characterized by confusion or delirium, often coupled with acute liver and kidney failure. Indeed, acute kidney injury is a common manifestation in individuals presenting with epidemic heatstroke. For example, in the 1995 heat wave in Chicago, over 50% of those presenting with heatstroke had acute kidney injury [9]. While acute kidney injury may accompany severe manifestations with coma and liver failure, milder forms of heatstroke may be only associated with fevers and acute kidney injury.

There appear to be 2 types of acute kidney injury [24]. One form appears to be classical rhabdomyolysis

(typically with creatine phosphokinase levels $>1,000$ μL), often associated with hyperuricemia and signs of dehydration. This form may be more common with exertional heatstroke. The other form is associated with normal or only mildly elevated creatine phosphokinase levels and is more common in epidemic heatstroke [24]. Indeed, unlike rhabdomyolysis, in which the injury appears more like an acute tubular injury, the second form of acute kidney injury clinically manifests more as an acute interstitial nephritis, with urinary leukocytosis and hematuria, and with a renal biopsy showing acute tubulointerstitial nephritis. It is thought that this condition results from ischemia, temperature-induced oxidative stress, and decreasing intracellular energy stores [30, 31].

Heatstroke is also commonly associated with electrolyte abnormalities [24, 32]. One study of 66 subjects with exertional heatstroke reported acute kidney injury in 91%, hyponatremia in 53%, hypokalemia in 71%, hypophosphatemia in 59%, hypocalcemia in 51%, and hypomagnesemia in 35% [32]. In particular, the low serum potassium, phosphate, and magnesium were all associated with increased urinary excretion of these electrolytes, suggesting a tubular defect. Other potential causes include loss of sodium and potassium through the sweat. Some subjects also present with respiratory alkalosis, which is known to reduce serum phosphate, although metabolic acidosis appears to be more common.

Some individuals (10–30%) with heatstroke-associated acute kidney injury require dialysis [32]. If the patient survives the acute illness, kidney function usually returns to normal [32]. However, some cases of heatstroke may progress to CKD months later with the presence of chronic tubulointerstitial nephritis on biopsy [33, 34].

Heat Stress Nephropathy as a Cause of CKD

In recent years, epidemics of CKD have been identified in various hot regions of the world where it preferentially affects workers who labor manually under extremely hot conditions [35]. One of the major sites of this disease is along the Pacific Coast of Central America, developing among sugarcane workers and others working in agricultural communities [36]. There is evidence that this epidemic has been progressively increasing since the 1970s [37]. The observation that the disease tends to occur in the hotter regions of Central America, coupled with evidence that the workers are placed under a great deal of heat stress [38, 39], has led to the hypothesis that the disease may be

driven by global warming [40]. Indeed, a recent study suggests that working in the sugarcane fields is associated with higher humidity due to the presence of the cane and that heat waves are driven not only by increasing mean temperatures but also by El Niño events [41].

There is increasing evidence that the development of CKD may result from repeated acute kidney injury driven by subclinical or clinical heatstroke [42]. Specifically, repeated acute kidney injury has been recently reported across work-shifts in sugarcane workers from this region [43–46]. While most cases are asymptomatic, some subjects present with fever, leukocytosis, leukocyturia, and acute kidney injury that may require admission to the local hospital [47–49]. These latter cases resemble heatstroke, as they may present with similar electrolyte abnormalities and also with acute interstitial nephritis on biopsy [47–49]. There is also evidence that some develop CKD over time [48], similar to that which occurs with exertional heatstroke [34].

Experimental studies support this association. Indeed recurrent heat stress and dehydration can induce chronic inflammation and tubular injury in mice and rats [50–52]. The mechanism of the kidney injury is likely related to increased internal body temperatures, the effects of hyperosmolarity to activate the polyol-fructokinase pathway, and the chronic effects of vasopressin to induce tubular and glomerular injury [50–52]. In addition, clinical studies suggest that the effects of heat and dehydration induce a concentrated and acidic urine, which can also lead to urinary urate crystallization with tubular damage [53]. Indeed, some experimental data suggest that lowering uric acid might provide protection [54, 55].

Acute kidney injury is now being reported throughout the world in hot agricultural communities including India (Andhra Pradesh), Sri Lanka (north central province), Mexico (Vera Cruz), central Florida, and the Central Valley of California [56–58]. In many of these areas, epidemics of CKD are also being reported [59–61]. A major concern is that these epidemics may be driven by increasing temperatures and heat waves, and that they may presage epidemics to come.

Other Effects of Heat Stress on the Kidney: Stones and Infections

Kidney stones (nephrolithiasis) are increasing in prevalence [62] and have also been proposed to result from increasing temperatures associated with climate change

[63]. Heat stress and dehydration predispose to urinary concentration and low urine volumes that increase the risk for stones [64]. In the United States, for example, the “stone belt” that characterizes the hotter regions in the southern United States is projected to move northward as climate warming continues [63]. Experimental studies show that the primary kidney stone substance associated with heat stress is uric acid, due to its increased generation following exercise-induced muscle damage and the urinary acidification that occurs during the concentrating process [64].

UTIs may also be related to underhydration and potentially affected by climate change. Indeed, a recent study found that increased daily water consumption could increase urine output and reduce the risk for UTIs [65].

Effect of Soft Drinks in Heat Stress-Associated Kidney Damage

Soft drinks contain fructose, a sugar that results in local tubular injury, inflammation, and oxidative stress when metabolized by the kidney [66]. Recent studies suggest that soft drinks may increase the risk for acute and chronic kidney injury [67]. Indeed, experimental studies have shown that rehydration with soft drinks could enhance kidney damage in dehydrated rats [51, 68]. In addition to the injury associated with fructose metabolism, fructose may be able to stimulate vasopressin that can then augment the renal injury [51, 69]. Indeed, a recent clinical study also reported that rehydration with soft drinks could induce markers of kidney damage in healthy subjects following exercise in high temperatures [70], although epidemiologic studies conducted in hot field settings to date have found no association.

Additive Effects of Toxins and Toxicants

A prevailing theory suggests that in the context of heat and dehydration, naturally occurring toxins and man-made toxicants may concentrate in the kidney during periods of recurrent acute kidney injury. Candidates under investigation include potentially nephrotoxic agrochemicals, heavy metals, use of nonsteroidal anti-inflammatory drugs, tobacco, and silica. Further research is ongoing, including environmental risk assessments that include meteorological conditions.

Summary

In summary, while the kidney has a major role in protecting the host from the effects of heat stress, it is also a target for heat stress associated injury. The effects of heat can lead to both acute and CKD, electrolyte abnormalities, and kidney stones and UTIs. As global warming continues, major efforts are required to assure adequate hydration and prevent overheating in vulnerable populations who are at risk for heatstroke. Heat warning systems, changes in occupational practices, and public health initiatives also are needed [71, 72]. Most importantly, scientific investigations should be directed at identifying how to slow, stop, and reverse global warming.

References

- 1 *Global Climate Change and Human Health*. San Francisco: Jossey-Bass; 2015.
- 2 Fischer EM, Knutti R. Anthropogenic contribution to global occurrence of heavy precipitation and high temperature extremes. *Nat Clim Chang*. 2015;5(6):560–4.
- 3 Meehl GA, Tebaldi C. More intense, more frequent, and longer lasting heat waves in the 21st century. *Science*. 2004 Aug;305(5686):994–7.
- 4 Rahmstorf S, Coumou D. Increase of extreme events in a warming world. *Proc Natl Acad Sci USA*. 2011 Nov;108(44):17905–9.
- 5 Luber G, McGeehin M. Climate change and extreme heat events. *Am J Prev Med*. 2008 Nov;35(5):429–35.
- 6 The Lancet. Heatwaves and health. *Lancet*. 2018 Aug;392(10145):359.
- 7 Watts N, Amann M, Ayeb-Karlsson S, Belesova K, Bouley T, Boykoff M, et al. The Lancet Countdown on health and climate change: from 25 years of inaction to a global transformation for public health. *Lancet*. 2018 Feb;391(10120):581–630.
- 8 Robine JM, Cheung SL, Le Roy S, Van Oyen H, Griffiths C, McNamee T, et al. Death toll exceeded 70,000 in Europe during the summer of 2003. *C R Biol*. 2008 Feb;331(2):171–8.
- 9 Dematte JE, O'Mara K, Buescher J, Whitney CG, Forsythe S, McNamee T, et al. Near-fatal heat stroke during the 1995 heat wave in Chicago. *Ann Intern Med*. 1998 Aug;129(3):173–81.
- 10 Mahant S. The evaluation and management of heat injuries in an intensive care unit. *Indian J Crit Care Med*. 2015 Aug;19(8):479–83.
- 11 Anonymous. India Heat Wave Kills More Than 1,400; Temperatures Soar To 118 Degrees, Associated Press, 2015. Available from:

Acknowledgments

This paper represents a contribution from the University of Colorado Climate Change and Health Consortium.

Disclosure Statement

R.J.J. has consulted for Danone Research, Astra Zeneca, Eli Lilly, and Horizon Pharmaceuticals and has equity with Colorado Research Partners LLC and XORT Therapeutics. He received travel expenses and registration fee from Danone Research to participate in the 2018 Hydration for Health Scientific Conference. L.G.S.-L., M.A.L., D.R.T., A.A.H., and C.A.R.-J. also have equity in Colorado Research Partners LLC. L.S.N. and J.B.-D. have received funding from Pantaleon for studies performed by the University of Colorado of workers in Guatemala. H.F.D., B.R.-I., Y.S., G.G., and J.L. report no conflicts of interest.

- https://weather.com/news/news/india-heat-wave-latest-news.
- 12 Dearden L. Karachi heat wave: Death toll tops 1,000 as government and electricity company trade blame: The Independent, 2015. Available from: <http://www.independent.co.uk/news/world/asia/pakistan-heat-wave-death-toll-tops-1000-as-government-and-electricity-company-trade-blame-10344719.html>.
- 13 Dally M, Butler-Dawson J, Krisher L, Monaghan A, Weitzenkamp D, Sorensen C, et al. The impact of heat and impaired kidney function on productivity of Guatemalan sugarcane workers. *PLoS One*. 2018 Oct;13(10):e0205181.
- 14 Friel S. Climate change, food insecurity and chronic diseases: sustainable and healthy policy opportunities for Australia. *N S W Public Health Bull*. 2010 May-Jun;21(5–6):129–33.
- 15 Kang S, Eltahir EA. North China Plain threatened by deadly heatwaves due to climate change and irrigation. *Nat Commun*. 2018 Jul;9(1):2894.
- 16 Pal JS, Eltahir EA. Future temperature in southwest Asia projected to exceed a threshold for human adaptability. *Nat Clim Chang*. 2016;6(2):197–200.
- 17 Im ES, Pal JS, Eltahir EA. Deadly heat waves projected in the densely populated agricultural regions of South Asia. *Sci Adv*. 2017 Aug;3(8):e1603322.
- 18 Vörösmarty CJ, Green P, Salisbury J, Lammers RB. Global water resources: vulnerability from climate change and population growth. *Science*. 2000 Jul;289(5477):284–8.
- 19 Kumm M, Ward PJ, de Moel H, Varis O. Is physical water scarcity a new phenomenon? Global assessment of water shortage over the last two millennia. *Environ Res Lett*. 2010;5:034006.
- 20 Morin C, Gandy J, Brazeilles R, Moreno LA, Kavouras SA, Martinez H, et al. Fluid intake patterns of children and adolescents: results of six Liq.In7 national cross-sectional surveys. *Eur J Nutr*. 2018 Jun;57(Suppl 3):113–23.
- 21 Borg M, Bi P, Nitschke M, Williams S, McDonald S. The impact of daily temperature on renal disease incidence: an ecological study. *Environ Health*. 2017 Oct;16(1):114.
- 22 Borg M, Nitschke M, Williams S, McDonald S, Nairn J, Bi P. Using the excess heat factor to indicate heatwave-related urinary disease: a case study in Adelaide, South Australia. *Int J Biometeorol*. 2019 Apr;63(4):435–47.
- 23 Tawatsupa B, Lim LL, Kjellstrom T, Seubsmann SA, Sleight A, Study Team TC; Thai Cohort Study Team. Association between occupational heat stress and kidney disease among 37,816 workers in the Thai Cohort Study (TCS). *J Epidemiol*. 2012;22(3):251–60.
- 24 Hart GR, Anderson RJ, Crumpler CP, Shulkin A, Reed G, Knochel JP. Epidemic classical heat stroke: clinical characteristics and course of 28 patients. *Medicine (Baltimore)*. 1982 May;61(3):189–97.
- 25 Adams T, Stacey E, Stacey S, Martin D. Exertional heat stroke. *Br J Hosp Med (Lond)*. 2012 Feb;73(2):72–8.
- 26 Bedno SA, Li Y, Han W, Cowan DN, Scott CT, Cavicchia MA, et al. Exertional heat illness among overweight U.S. Army recruits in basic training. *Aviat Space Environ Med*. 2010 Feb;81(2):107–11.
- 27 Casa DJ, Armstrong LE, Ganio MS, Yeargin SW. Exertional heat stroke in competitive athletes. *Curr Sports Med Rep*. 2005 Dec;4(6):309–17.
- 28 Goforth CW, Kazman JB. Exertional heat stroke in navy and marine personnel: a hot topic. *Crit Care Nurse*. 2015 Feb;35(1):52–9.

- 29 Staal Wåsterlund D. Managing heat in agricultural work: increasing worker safety and productivity by controlling heat exposure. Forestry Working Paper. Rome, Food and Agricultural Organization of the United Nations. 2018;1:1–67.
- 30 Zager RA, Altschuld R. Body temperature: an important determinant of severity of ischemic renal injury. *Am J Physiol*. 1986 Jul;251(1 Pt 2):F87–93.
- 31 Zager RA. Hyperthermia: effects on renal ischemic/reperfusion injury in the rat. *Lab Invest*. 1990 Sep;63(3):360–9.
- 32 Satirapoj B, Kongthaworn S, Choovichian P, Supasyndh O. Electrolyte disturbances and risk factors of acute kidney injury patients receiving dialysis in exertional heat stroke. *BMC Nephrol*. 2016 Jun;17(1):55.
- 33 Kew MC, Abrahams C, Levin NW, Seftel HC, Rubenstein AH, Bersohn I. The effects of heatstroke on the function and structure of the kidney. *Q J Med*. 1967 Jul;36(143):277–300.
- 34 Kew MC, Abrahams C, Seftel HC. Chronic interstitial nephritis as a consequence of heatstroke. *Q J Med*. 1970 Apr;39(154):189–99.
- 35 Johnson RJ, Wesseling C, Newman LS. Epidemic Chronic Kidney Disease in Agricultural Communities. *N Engl J Med*. Forthcoming 2019.
- 36 Correa-Rotter R, Wesseling C, Johnson RJ. CKD of unknown origin in Central America: the case for a Mesoamerican nephropathy. *Am J Kidney Dis*. 2014 Mar;63(3):506–20.
- 37 Wesseling C, van Wendel de Joode B, Crowe J, Rittner R, Sanati NA, Hogstedt C, et al. Mesoamerican nephropathy: geographical distribution and time trends of chronic kidney disease mortality between 1970 and 2012 in Costa Rica. *Occup Environ Med*. 2015 Oct;72(10):714–21.
- 38 Crowe J, Nilsson M, Kjellstrom T, Wesseling C. Heat-related symptoms in sugarcane harvesters. *Am J Ind Med*. 2015 May;58(5):541–8.
- 39 Crowe J, Wesseling C, Solano BR, Umaña MP, Ramírez AR, Kjellstrom T, et al. Heat exposure in sugarcane harvesters in Costa Rica. *Am J Ind Med*. 2013 Oct;56(10):1157–64.
- 40 Glaser J, Lemery J, Rajagopalan B, Diaz HF, García-Trabanino R, Taduri G, et al. Climate Change and the Emergent Epidemic of CKD from Heat Stress in Rural Communities: The Case for Heat Stress Nephropathy. *Clin J Am Soc Nephrol*. 2016 Aug;11(8):1472–83.
- 41 Diaz HF, Mora C, Wesseling C, Johnson RJ, Crowe J, Hidalgo HG, et al. Increasing Heat Stress, Kidney Disease, and Possible Connection to Climate Change in Selected Regions of Central America. *Clim Change*. Forthcoming 2019.
- 42 Johnson RJ. Pro: Heat stress as a potential etiology of Mesoamerican and Sri Lankan nephropathy: a late night consult with Sherlock Holmes. *Nephrol Dial Transplant*. 2017 Apr;32(4):598–602.
- 43 García-Trabanino R, Jarquín E, Wesseling C, Johnson RJ, González-Quiroz M, Weiss I, et al. Heat stress, dehydration, and kidney function in sugarcane cutters in El Salvador—A cross-shift study of workers at risk of Mesoamerican nephropathy. *Environ Res*. 2015 Oct;142:746–55.
- 44 Wesseling C, Aragón A, González M, Weiss I, Glaser J, Rivard CJ, et al. Heat stress, hydration and uric acid: a cross-sectional study in workers of three occupations in a hotspot of Mesoamerican nephropathy in Nicaragua. *BMJ Open*. 2016 Dec;6(12):e011034.
- 45 Solis Zepeda GA. Impacto de las medidas preventivas para evitar el deterioro de la función renal por el Síndrome de Golpe por Calor en trabajadores agrícolas del Ingenio San Antonio del Occidente de Nicaragua, Ciclo Agrícola 2005–2006. Internal Medicine Department. León: Universidad Nacional Autónoma de Nicaragua; 2007.
- 46 Sorensen CJ, Butler-Dawson J, Dally M, Krisher L, Griffin BR, Johnson RJ, et al. Risk Factors and Mechanisms Underlying Cross-shift Decline in Kidney Function in Guatemalan Sugarcane Workers. *J Occup Environ Med*. 2019 Mar;61(3):239–50.
- 47 Fischer RS, Mandayam S, Chavarria D, Vangala C, Nolan MS, Garcia LL, et al. Clinical Evidence of Acute Mesoamerican Nephropathy. *Am J Trop Med Hyg*. 2017 Oct;97(4):1247–56.
- 48 Fischer RS, Vangala C, Mandayam S, Chavarria D, García-Trabanino R, Garcia F, et al. Clinical markers to predict progression from acute to chronic kidney disease in Mesoamerican nephropathy. *Kidney Int*. 2018 Dec;94(6):1205–16.
- 49 Fischer RS, Vangala C, Truong L, Mandayam S, Chavarria D, Granera Llanes OM, et al. Early detection of acute tubulointerstitial nephritis in the genesis of Mesoamerican nephropathy. *Kidney Int*. 2018 Mar;93(3):681–90.
- 50 Roncal Jimenez CA, Ishimoto T, Lanaspá MA, Rivard CJ, Nakagawa T, Ejaz AA, et al. Fructokinase activity mediates dehydration-induced renal injury. *Kidney Int*. 2014 Aug;86(2):294–302.
- 51 García-Arroyo FE, Cristóbal M, Arellano-Buendía AS, Osorio H, Tapia E, Soto V, et al. Rehydration with soft drink-like beverages exacerbates dehydration and worsens dehydration-associated renal injury. *Am J Physiol Regul Integr Comp Physiol*. 2016 Jul;311(1):R57–65.
- 52 García-Arroyo FE, Tapia E, Blas-Marrón MG, Gonzaga G, Silverio O, Cristóbal M, et al. Vasopressin Mediates the Renal Damage Induced by Limited Fructose Rehydration in Recurrently Dehydrated Rats. *Int J Biol Sci*. 2017 Jul;13(8):961–75.
- 53 Roncal-Jimenez C, García-Trabanino R, Barregard L, Lanaspá MA, Wesseling C, Harra T, et al. Heat Stress Nephropathy From Exercise-Induced Uric Acid Crystalluria: A Perspective on Mesoamerican Nephropathy. *Am J Kidney Dis*. 2016 Jan;67(1):20–30.
- 54 Roncal-Jimenez CA, Sato Y, Milagres T, Andres Hernandez A, García G, Bjornstad P, et al. Experimental heat stress nephropathy and liver injury are improved by allopurinol. *Am J Physiol Renal Physiol*. 2018 Sep;315(3):F726–33.
- 55 Sánchez-Lozada LG, García-Arroyo FE, Gonzaga G, Silverio O, Blas-Marrón MG, Muñoz-Jimenez I, et al. Kidney Injury from Recurrent Heat Stress and Rhabdomyolysis: Protective Role of Allopurinol and Sodium Bicarbonate. *Am J Nephrol*. 2018;48(5):339–48.
- 56 Mix J, Elon L, Vi Thien Mac V, Flocks J, Economos E, Tovar-Aguilar AJ, et al. Hydration Status, Kidney Function, and Kidney Injury in Florida Agricultural Workers. *J Occup Environ Med*. 2018 May;60(5):e253–60.
- 57 Moyce S, Mitchell D, Armitage T, Tancredi D, Joseph J, Schenker M. Heat strain, volume depletion and kidney function in California agricultural workers. *Occup Environ Med*. 2017 Jun;74(6):402–9.
- 58 Badurdeen Z, Nanayakkara N, Ratnatunga NV, Wazil AW, Abeysekera TD, Rajakrishna PN, et al. Chronic kidney disease of uncertain etiology in Sri Lanka is a possible sequel of interstitial nephritis! *Clin Nephrol*. 2016;86(13):106–9.
- 59 Wijetunge S, Ratnatunga NV, Abeysekera TD, Wazil AW, Selvarajah M. Endemic chronic kidney disease of unknown etiology in Sri Lanka: correlation of pathology with clinical stages. *Indian J Nephrol*. 2015 Sep-Oct;25(5):274–80.
- 60 Mendoza-González MF, Montes-Villaseñor E, Muñoz-Flores P, Salado-Pérez M, Espejo-Guevara DM, Tapia-Jaime G. Prevalencia de Enfermedad Renal Crónica en una Población de Alto Riesgo. Tierra Blanca, Veracruz, México: Memorias Convención Internacional de Salud Pública Havana, Cuba, Cuba Salud; 2012.
- 61 Ganguli A. Uddanam Nephropathy/Regional Nephropathy in India: Preliminary Findings and a Plea for Further Research. *Am J Kidney Dis*. 2016 Sep;68(3):344–8.
- 62 Stamatelou KK, Francis ME, Jones CA, Nyberg LM Jr, Curhan GC. Time trends in reported prevalence of kidney stones in the United States: 1976–1994. *Kidney Int*. 2003 May;63(5):1817–23.
- 63 Brikowski TH, Lotan Y, Pearle MS. Climate-related increase in the prevalence of urolithiasis in the United States. *Proc Natl Acad Sci USA*. 2008 Jul;105(28):9841–6.
- 64 Borghi L, Meschi T, Amato F, Novarini A, Romanelli A, Cigala F. Hot occupation and nephrolithiasis. *J Urol*. 1993 Dec;150(6):1757–60.

- 65 Hooton TM, Vecchio M, Iroz A, Tack I, Dornic Q, Seksek I, et al. Effect of Increased Daily Water Intake in Premenopausal Women With Recurrent Urinary Tract Infections: A Randomized Clinical Trial. *JAMA Intern Med.* 2018 Nov;178(11):1509–15.
- 66 Cirillo P, Gersch MS, Mu W, Scherer PM, Kim KM, Gesualdo L, et al. Ketohexokinase-dependent metabolism of fructose induces proinflammatory mediators in proximal tubular cells. *J Am Soc Nephrol.* 2009 Mar;20(3):545–53.
- 67 Shoham DA, Durazo-Arvizu R, Kramer H, Luke A, Vupputuri S, Kshirsagar A, et al. Sugary soda consumption and albuminuria: results from the National Health and Nutrition Examination Survey, 1999-2004. *PLoS One.* 2008;3(10):e3431.
- 68 Milagres T, García-Arroyo FE, Lanaspá MA, García G, Ishimoto T, Andres-Hernando A, et al. Rehydration with fructose worsens dehydration-induced renal damage. *BMC Nephrol.* 2018 Jul;19(1):180.
- 69 Roncal-Jimenez CA, Milagres T, Andres-Hernando A, Kuwabara M, Jensen T, Song Z, et al. Effects of exogenous desmopressin on a model of heat stress nephropathy in mice. *Am J Physiol Renal Physiol.* 2017 Mar;312(3):F418–26.
- 70 Chapman CL, Johnson BD, Sackett JR, Parker MD, Schlader ZJ. Soft drink consumption during and following exercise in the heat elevates biomarkers of acute kidney injury. *Am J Physiol Regul Integr Comp Physiol.* 2019 Mar;316(3):R189–98.
- 71 Toloo GS, Fitzgerald G, Aitken P, Verrall K, Tong S. Are heat warning systems effective? *Environ Health.* 2013 Apr;12(1):27.
- 72 Mayrhuber EA, Dücker ML, Wallner P, Arnerberger A, Alex B, Wiesböck L, et al. Vulnerability to heatwaves and implications for public health interventions – A scoping review. *Environ Res.* 2018 Oct;166:42–54.